

chain altered distribution in a much higher proportion of patients with X-linked AS [5].

It is our experience that skin biopsy, examined with conventional, and if needed, with confocal microscopy, is able to virtually detect (almost) all cases of X-linked AS [5], thus allowing to avoid or to postpone more invasive and/or expensive diagnostic procedure like renal biopsy and genetic investigations, which, in addition, are not diagnostic in all cases.

GIANFRANCO RIZZONI and LAURA MASSELLA
Rome, Italy

Correspondence to Laura Massella, Division of Nephrology, Bambino Gesù Children's Hospital and Research Institute, Piazza S. Onofrio 14, 00165 Rome, Italy.
E-mail: lmassella@opbg.net

REFERENCES

1. GREGORY MC: Alport syndrome and thin basement membrane nephropathy: Unraveling the tangled strands of type IV collagen. *Kidney Int* 65:1109–1110, 2004
2. KASHTAN CE: Alport syndrome and thin glomerular basement membrane disease. *J Am Soc Nephrol* 9:1736–1750, 1998
3. RIZZONI G, MASSELLA L, MUDA AO: Alpha 5 COLIV chain distribution in glomerular basement membrane in a male with X-linked Alport syndrome and thin basement membrane. *Pediatr Nephrol* 15:325, 2000
4. KASHTAN CE: Uptodate, 2003
5. ONETTI MUDA A, MASSELLA L, GIANNAKAKIS K, *et al*: Confocal microscopy of the skin in the diagnosis of X-linked Alport syndrome. *J Invest Dermatol* 121:208–211, 2003

Reply from the Author

I thank Professors Kashtan and Rizzoni and Dr. Massella for their comments. They rightly point out the value of immunostaining of renal biopsies for the diagnosis of X-linked and autosomal Alport syndrome and the utility of skin immunofluorescence in diagnosis of many cases of X-linked Alport syndrome. The use of confocal laser scanning microscopy (CLSM) [1–3] is an elegant technique to improve the spatial resolution of $\alpha 5(\text{IV})$ chain distribution in basement membranes.

Ueda *et al* [3], also using CLSM, have reported that compared with $\alpha 2(\text{IV})$, $\alpha 5(\text{IV})$ expression in GBM is reduced in patients with TBMD. The reduction in $\alpha 5(\text{IV})$ but not $\alpha 3(\text{IV})$ and $\alpha 4(\text{IV})$ is difficult to reconcile with the genetic evidence implicating mutations of COL4A3 and COL4A4 in TBMD.

Confirmation and clarification of the results of CLSM of EBM and GBM is eagerly awaited. Until sequencing of the relevant collagen genes or other comprehensive genetic testing is routinely and readily available, immunofluorescence of skin biopsies is a minimally invasive means for diagnosis of many cases of X-linked Alport syndrome. If quantitative comparisons of the different $\alpha(\text{IV})$ chains in GBM permit a positive diagnosis of TBMD, this will further strengthen our diagnostic hand.

MARTIN GREGORY
Salt Lake City, Utah

Correspondence to Martin Gregory, Department of Medicine, University of Utah Health Sciences Center, Salt Lake City, UT.
E-mail: martin.gregory@hsc.utah.edu

REFERENCES

1. MUDA AO, MASSELLA L, GIANNAKAKIS K, *et al*: Confocal microscopy of the skin in the diagnosis of X-linked Alport syndrome. *J Invest Dermatol* 121:208–211, 2003
2. MASSELLA L, ONETTI MUDA A, FARAGGIANA T, *et al*: Epidermal basement membrane alpha 5(IV) expression in females with Alport syndrome and severity of renal disease. *Kidney Int* 64:1787–1791, 2003
3. UEDA T, NAKAJIMA M, AKAZAWA H, *et al*: Quantitative analysis of glomerular type IV collagen alpha3–5 chain expression in children with thin basement membrane disease. *Nephron* 92:271–278, 2002

Diagnosis of Alport syndrome

To the Editor: Professor Gregory's recent editorial [1] addresses the difficulties that may arise in attempts to distinguish Alport syndrome and thin basement membrane nephropathy, particularly in children. Positive diagnosis of the X-linked and autosomal-recessive forms of Alport syndrome may be made by immunostaining of renal biopsy specimens for alpha chains of type IV collagen [2, 3]. X-linked Alport syndrome can also be diagnosed by immunostaining of skin biopsy specimens for the alpha5 chain of type IV collagen [4].

There are patients who exhibit expression of type IV collagen chains that is indistinguishable from control, despite the presence of pathogenic mutations in the COL4A3, COL4A4, or COL4A5 gene [3, 4]. In these cases, estimation of prognosis and genetic counseling would be greatly aided by the availability of molecular genetic diagnostic services. More than 80% of mutations in the COL4A5 gene can be identified by direct nucleotide sequencing [5].

CLIFFORD E. KASHTAN
Minneapolis, Minnesota

Correspondence to Clifford E. Kashtan, University of Minnesota, MMC 491, 420 Delaware St. SE, Minneapolis, MN 55455.
E-mail: kasht001@maroon.tc.umn.edu

REFERENCES

1. GREGORY MC: Alport syndrome and thin basement membrane nephropathy: Unraveling the tangled strands of type IV collagen. *Kidney Int* 65:1109–1110, 2004
2. NAKANISHI K, YOSHIKAWA N, IJIMA K, *et al*: Immunohistochemical study of $\alpha 1$ –5 chains of type IV collagen in hereditary nephritis. *Kidney Int* 46:1413–1421, 1994
3. GUBLER MC, KNEBELMANN B, BEZIAU A, *et al*: Autosomal recessive Alport syndrome: Immunohistochemical study of type IV collagen chain distribution. *Kidney Int* 47:1142–1147, 1995
4. VAN DER LOOP FTL, MONNENS LAH, SCHRODER CH, *et al*: Identification of COL4A5 defects in Alport syndrome by immunochemistry of skin. *Kidney Int* 55:1217–1224, 1999

5. MARTIN P, HEISKARI N, ZHOU J, *et al*: High mutation detection rate in the COL4A5 collagen gene in suspected Alport syndrome using PCR and direct DNA sequencing. *J Am Soc Nephrol* 9:2291–2301, 1998

Reply from the Author

I thank Professors Kashtan and Rizzoni and Dr. Massella for their comments. They rightly point out the value of immunostaining of renal biopsies for the diagnosis of X-linked and autosomal Alport syndrome and the utility of skin immunofluorescence in diagnosis of many cases of X-linked Alport syndrome. The use of confocal laser scanning microscopy (CLSM) [1–3] is an elegant technique to improve the spatial resolution of $\alpha 5(\text{IV})$ chain distribution in basement membranes.

Ueda *et al* [3], also using CLSM, have reported that compared with $\alpha 2(\text{IV})$, $\alpha 5(\text{IV})$ expression in GBM is reduced in patients with TBMD. The reduction in $\alpha 5(\text{IV})$ but not $\alpha 3(\text{IV})$ and $\alpha 4(\text{IV})$ is difficult to reconcile with the genetic evidence implicating mutations of COL4A3 and COL4A4 in TBMD.

Confirmation and clarification of the results of CLSM of EBM and GBM is eagerly awaited. Until sequencing of the relevant collagen genes or other comprehensive genetic testing is routinely and readily available, immunofluorescence of skin biopsies is a minimally invasive means for diagnosis of many cases of X-linked Alport syndrome. If quantitative comparisons of the different $\alpha(\text{IV})$ chains in GBM permit a positive diagnosis of TBMD, this will further strengthen our diagnostic hand.

MARTIN GREGORY
Salt Lake City, Utah

Correspondence to Martin Gregory, Department of Medicine, University of Utah Health Sciences Center, Salt Lake City, UT.
E-mail: martin.gregory@hsc.utah.edu

REFERENCES

1. MUDA AO, MASSELLA L, GIANNAKAKIS K, *et al*: Confocal microscopy of the skin in the diagnosis of X-linked Alport syndrome. *J Invest Dermatol* 121:208–211, 2003
2. MASSELLA L, ONETTI MUDA A, FARAGGIANA T, *et al*: Epidermal basement membrane $\alpha 5(\text{IV})$ expression in females with Alport syndrome and severity of renal disease. *Kidney Int* 64:1787–1791, 2003
3. UEDA T, NAKAJIMA M, AKAZAWA H, *et al*: Quantitative analysis of glomerular type IV collagen $\alpha 3\text{--}5$ chain expression in children with thin basement membrane disease. *Nephron* 92:271–278, 2002

Dialysis dose and gender: A different hypothesis

To the Editor: A recent report [1] suggested that females who are smaller than males need higher dialysis




Fig. 1. Illustration of three relationships discriminating the high (upper symbols) and low (lower symbols) Kt/V groups for males (solid circles) and females (open squares) in the HEMO study. The relationships are 1) $Kt = 1.16 V$ (thin black line) representing the low dose HEMO study Kt/V for both genders, 2) $Kt = 30 + 0.5 V$ (thick black line), and 3) $35 + 0.3 V$ (thick gray line).

dose (Kt/V) than males. The idea that smaller persons require higher Kt per L of V is not new [2], and suggests that a 0-intercept rule for judging Kt per unit of V is incorrect [3].

Figure 1 illustrates a non-0 intercept rule (thick black line: $Kt = 30 + 0.5 V$) [3]. The mean $Kt \pm SD$ (Table 3) [1] is shown for females (squares) and males (circles) at the mean V (Table 2) [1] for the high (upper symbols) and low (lower symbols) hemodialysis (HEMO) treatment groups. The steep, thin black line is a 0-intercept Kt/V rule ($Kt = 0 + 1.16 V$) [1]. All groups were treated at or above the Kt for their V by that rule. Females in the low treatment group ($Kt = 38.2$; $eKt/V = 1.17$), however, had marginally worse ($P = 0.02$) survival than females in the high treatment group [1].

The gray line ($Kt = 35 + 0.3 V$) is a rotation of the earlier black line [3], and better discriminates among groups according to outcome. Only low Kt/V females received substantially suboptimal therapy according to this illustrative rule.

The point of this exercise was not to recommend a new rule for judging treatment. It only shows that one need not resort to speculations about different uremic toxin generation in males and females [1] to explain this marginal survival difference. All one needs to do is accept the possibility that a 0-intercept Kt/V rule may be suboptimal.

EDMUND G. LOWRIE, NORMA OFSTHUN, and ZHENGSHENG LI
Lexington, Massachusetts

Correspondence to Edmund G. Lowrie, 21 Edmonds Road, Concord, MA 01742.
E-mail: edlowrie@prodigy.net

REFERENCES

1. DEPNER T, DAUGIRDAS J, GREENE T, *et al*: Dialysis dose and the effect of gender and body size on outcome in the HEMO study. *Kidney Int* 65:1386–1394, 2004